

A Comment about the existence of a weak solution for a nonlinear Wave Damped Propagation

LUIZ C.L. BOTELHO

Departamento de Matemática Aplicada,
Instituto de Matemática, Universidade Federal Fluminense,
Rua Mario Santos Braga
24220-140, Niterói, Rio de Janeiro, Brazil
e-mail: botelho.luiz@superig.com.br

Abstract

We give a proof for the existence of weak solutions on the initial-value problem of a non-linear wave damped propagation.

1 Introduction

One of the most important problems in the mathematical physics of the non-linear diffusion and wave-damped propagation is to establish the existence and the uniqueness of weak solutions in some convenient Banach spaces for the associated nonlinear evolution equation (see Refs 1–5). Another important class is these coming from initial-value problems for non-linear diffusion or damped hyperbolic partial differential equations with random initial conditions associated to the Gaussian processes sampled in certain Hilbert Spaces ([2]) and simulating the statistical turbulence physical phenomena ([6]).

The purpose of this short note is to contribute to such mathematical physics studies by using functional spaces compacity argument in order to produce proofs for the existence of weak solutions for a class of non-linear wave damped propagation on a smooth domain with Dirichlet boundary conditions and initial values belonging to the space $L^2(\Omega)$.

2 Existence Solution for Hyperbolic Non-linear Damping

Let us introduce firstly a non-linearity associated to a real valued function $F(x)$ on R such that $F'(x)$ is strictly positive and an external forcing source $f(x, t)$ supposed to be on the space $L^\infty([0, T], L^2(\Omega))$.

Associated to them we consider the following non-linear damped Hyperbolic initial value problem on $\Omega \times [0, T]$ with imposed Dirichlet boundary conditions and a damping positive constant ν .

$$\frac{\partial^2 U(x, t)}{\partial t^2} + (AU)(x, t) = -\nu \frac{\partial U(x, t)}{\partial t} + \Delta \left(F \left(\frac{\partial U}{\partial t}(x, t) \right) \right) + f(x, t) \quad (1)$$

The $L^2(\Omega)$ -initial conditions are given by

$$U(x, 0) = g(x) \in L^2(\Omega) \quad (2-a)$$

$$U_t(x, 0) = h(x) \in L^2(\Omega) \quad (2-b)$$

$$U(x, t)|_{\partial\Omega} = 0 \quad (2-c)$$

and now the non-homogenous term $f(x, t)$ is considered to be a function belonging to the functional space

$$L^2([0, T] \times \Omega) \cap L^\infty([0, T], L^2(\Omega)). \quad (3)$$

We have thus the following theorem of existence (without uniqueness)

Theorem 2. *There exists a solution $\bar{U}(x, t)$ on $L^\infty([0, T] \times L^2(\Omega))$ for eq.(1)–eq.(3) in the weak sense with a test functional space as given by $C_0^\infty([0, T], H^2(\Omega) \cap H_0^1(\Omega))$.*

In order to arrive at such theorem, let us consider the a priori estimate for eq.(1) with $\dot{U}^{(n)} \equiv \frac{\partial}{\partial t}(U^{(n)}(x, t))$. Namely:

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|\dot{U}^{(n)}\|_{L^2(\Omega)}^2 + \nu \|\dot{U}^{(n)}\|_{L^2(\Omega)}^2 \\ & + (A\dot{U}^{(n)}, U^{(n)})_{L^2(\Omega)} + \|(F'(\dot{U}^{(n)}))^{1/2} \nabla \dot{U}^{(n)}\|_{L^2(\Omega)}^2 \\ & = (f, \dot{U}^{(n)})_{L^2(\Omega)} \end{aligned} \quad (4)$$

or equivalently

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \left\{ \|\dot{U}^{(n)}\|_{L^2(\Omega)}^2 + (AU^{(n)}, U^{(n)})_{L^2(\Omega)} \right\} \\
& + \nu \left\{ \|\dot{U}^{(n)}\|_{L^2(\Omega)}^2 + (AU^{(n)}, U^{(n)})_{L^2(\Omega)} \right\} \\
& + \|(F'(\dot{U}^{(n)}))^{1/2} \nabla \dot{U}^{(n)}\|_{L^2(\Omega)}^2 \leq \nu (AU^{(n)}, U^{(n)}) \\
& + \frac{1}{2} \left(p \|f\|_{L^2(\Omega)}^2 + \frac{1}{p} \|\dot{U}^{(n)}\|_{L^2(\Omega)}^2 \right)
\end{aligned} \tag{5}$$

If one chooses here the integer p such that $\frac{1}{2p} = \nu$, we have the simple bound

$$\frac{1}{2} \frac{d}{dt} \left\{ \|\dot{U}^{(n)}\|_{L^2(\Omega)}^2 + (AU^{(n)}, U^{(n)})_{L^2(\Omega)} \right\} \leq \frac{1}{4\nu} \|f\|_{L^2(\Omega)}^2. \tag{6}$$

As a consequence of eq.(6), there is a constant M such that the uniform bounds holds true (even if for the case $T = +\infty$ for the case of $f \in L^2([0, \infty), L^2(\Omega))$).

$$\sup \text{ess}_{0 \leq t \leq T} \|\dot{U}^{(n)}\|_{L^2(\Omega)}^2 \leq M \tag{7}$$

$$\sup \text{ess}_{0 \leq t \leq T} \|U^{(n)}\|_{L^2(\Omega)}^2 \leq M. \tag{8}$$

As a consequence of the bounds eq.(7) – eq.(8), there are two functions $\overline{U}(x, t)$ and $\overline{P}(x, t)$ such that we have the weak star convergence on $L^\infty([0, T] \times L^2(\Omega))$

$$\text{weak-star} \quad \lim_{n \rightarrow \infty} (U^{(n)}(x, t)) = \overline{U}(x, t) \tag{9}$$

$$\text{weak-star} \quad \lim_{n \rightarrow \infty} \left(\frac{\partial}{\partial t} U^{(n)}(x, t) \right) = \overline{P}(x, t). \tag{10}$$

We have thus that the relationship below hold true for any test function $v(x, t) \in C_c^\infty([0, T] \times H^2(\Omega) \cap H_0^1(\Omega))$ obviously satisfying the relations $v(x, 0) = v(x, T) = \Delta v(x, 0) = \Delta v(x, T) = v_t(x, 0) = v_t(x, T) = v_{tt}(x, 0) = v_{tt}(x, T) \equiv 0$ as a consequence of applying the Aubin-Lion theorem

$$\begin{aligned}
& \int_0^T dt \left[\left(\overline{U}, \frac{d^2 v}{dt^2} \right)_{L^2(\Omega)} + (\overline{U}, Av)_{L^2(\Omega)} \right. \\
& \quad \left. + \nu \left(\overline{U}, -\frac{dv}{dt} \right)_{L^2(\Omega)} - (F(\overline{P}), \Delta v)_{L^2(\Omega)} \right] \\
& = \int_0^T dt (f, v)_{L^2(\Omega)}
\end{aligned} \tag{11}$$

with the initial conditions

$$\overline{U}(x, 0) = g(x) \in L^2(\Omega) \quad (12)$$

$$\overline{P}(x, 0) = h(x) \in L^2(\Omega). \quad (13)$$

Let us show now that

$$\overline{U}(x, t) = \int_0^t ds \overline{P}(x, s). \quad (14)$$

Firstly, let us remark that integrating on the interval $0 \leq t \leq T$ the relationship eq.(4), one obtains the following estimate

$$\begin{aligned} & \frac{1}{2} \left(\|\dot{U}^n(t)\|_{L^2(\Omega)}^2 - \|\dot{U}^n(0)\|_{L^2(\Omega)}^2 \right) \\ & + \left(\nu \int_0^t ds \|\dot{U}_n\|_{L^2(\Omega)}^2(s) \right) + \frac{1}{2} (AU^{(n)}(t), U^{(n)}(t))_{L^2(\Omega)} \\ & - \frac{1}{2} (AU^{(n)}(0), U^{(n)}(0)) + \int_0^t ds \|(F'(\dot{U}^{(n)}))^{1/2} \nabla \dot{U}^{(n)}\|_{L^2(\Omega)}^2 \\ & \leq \frac{p'}{2} \int_0^t ds \|f\|_{L^2(\Omega)}^2 + \frac{1}{2p'} \int_0^t \|\dot{U}_n(s)\|^2 ds \end{aligned} \quad (15)$$

Since the operator A satisfies the Gårding-Poincaré inequality on $L^2(\Omega)$

$$(AU^{(n)}, U^{(n)})_{L^2(\Omega)}(t) \geq \gamma(\Omega) \|U^{(n)}\|_{L^2(\Omega)}^2(t), \quad (16)$$

one can see straightforwardly from eq.(15) by choosing $2p' > \frac{1}{\nu}$ and the previous bounds eq.(7) that there is a positive constant B such that

$$\int_0^T ds \|\dot{U}^{(n)}(s)\|_{L^2(\Omega)}^2 \leq B = MT \quad (17)$$

which by its turn yields that $\frac{dU_n(t,x)}{dt}$ is weakly convergent to $\overline{P}(x, t)$ in $L^2([0, T], L^2(\Omega))$.

As a consequence of general theorems of Function convergence on space of integrable functions (Aubin-Lion theorem again ([1])), one has that $\overline{P}(x, t)$ is the time-derivative of the function $\overline{U}(x, t)$ almost everywhere on Ω , since it is expected that $U^{(n)}(x, t)$ should be a strongly convergent sequence to $\overline{U}(x, t)$ on the separable and reflexive Banach Space $L^\infty([0, T] \times L^2(\Omega))$ ([1]). (See Appendix A for mathematical details).

APPENDIX A

Let us show that the functions $\overline{U}(x, t) \in L^\infty([0, T], L^2(\Omega))$ given by eq.(8) and $\overline{P}(x, t) \in L^\infty([0, T], L^2(\Omega))$ – eq.(9) are *coincident as elements* of the above written functional space of $L^2(\Omega)$ – valued essential bounded functions on $(0, T)$.

To verify such result, let us call the reader attention that since $U_m(x, t)$ is weakly convergent to $U(x, t)$ in $L^\infty([0, T], L^2(\Omega))$, we have that the set $\{U_m(x, t)\}$ is convergent to the function $U(x, t)$ as a Schwartz distribution on $L^2(\Omega)$ since $D([0, T], L^2(\Omega)) \subset L^1([0, T], L^2(\Omega))$.

This means that

$$U_m(x, t) \rightarrow U(x, t) \quad \text{in} \quad D'([0, T], L^2(\Omega)). \quad (1-A)$$

Analogous result hold true for the time-derivative of the above written equation as a result of $U(x, t)$ be a function

$$\frac{\partial}{\partial t} U_m(x, t) \rightarrow \frac{\partial}{\partial t} U(x, t) \quad \text{in} \quad D'([0, T], L^2(\Omega)) \quad (2-A)$$

By the other side, the set $\left\{ \frac{\partial U_m(x, t)}{\partial t} \right\}$ converges weakly star in $L^1([0, T], L^2(\Omega))$ to $\overline{P}(x, t) \in L^\infty([0, T], L^2(\Omega))$ which, by its turn, means that

$$\frac{\partial U_m(x, t)}{\partial t} \rightarrow \overline{P}(x, t) \quad \text{in} \quad D'([0, T], L^2(\Omega)). \quad (3-A)$$

By the uniqueness of the limit on the distributional space $D'([0, T], L^2(\Omega))$, we have the coincidence of $\frac{\partial U(x, t)}{\partial t}$ and $\overline{P}(x, t)$ as elements of $D'([0, T], L^2(\Omega))$. However, $\overline{P}(x, t)$ is a function, so by general theorems on Schwartz distribution theory $\left\{ \frac{\partial U(x, t)}{\partial t} \right\}$ must be a function either since $L^2(\Omega)$ is a separable Hilbert Space. As a consequence we have that $\frac{\partial U(x, t)}{\partial t} = \overline{P}(x, t)$ as elements of $L^\infty([0, T], L^2(\Omega))$, which is the result searched

$$\frac{\partial U(x, t)}{\partial t} = \overline{P}(x, t) \quad \text{a.e. in} \quad ([0, T] \times \Omega). \quad (4-C)$$

References:

- [1] J. Wloka, “Partial differential equations”, Cambridge University Press, (1987).
- [2] Y. Yamasaki – “Measures on Infinite Dimensional Spaces”, Series in Pure Mathematics World Scientific - vol. 5, (1985).
 - B. Simon, Functional Integration and Quantum Physics, Academic Press, (1979).
 - J. Glimm and A. Jaffe, “Quantum Physics – a functional Integral point of view” – Springer Verlag, (1981).
 - Xia Dao Xing – “Measure and Integration Theory on Infinite Dimensional Space”, Academic Press, (1972).
 - L. Schwartz, “Random Measures on Arbitrary Topological Space and Cylindrical Measures - Tata Institute - Oxford University Press, (1973).
 - Ya G. Sinai, “The theory of phase transitions: rigorous results, London, Pergamon Press, (1981).
- [3] O. A. Ladyzenskaja, V.A. Solonnikov and N.N. Ural'ceva - “Linear and Quasilinear Equations of Parabolic Type”, Amer. Math. Soc. Transl., A.M.S., Providence, R.I., (1968).
- [4] S.G. Mikhlin – “Mathematical Physics, an advanced course”, Series in Applied Mathematics and Mechanics, North-Holland, (1970).
- [5] Luiz C.L. Botelho – Journal of Physics A: Math. Gen., 34, L 131–L 137, (2001); Modern Phys. Lett. 16B, No. 21, 793–8-6, (2002).
 - A. Bensoussan and R. Teman, Journal of Functional Analysis 13, 195–222, (1973).
 - Avner Friedman – “Variational Principles and Free-Boundary Problems” – Pure & Applied Mathematics, John-Wiley & Sony, N.Y., (1982).
- [6] Luiz C.L. Botelho – Il Nuovo Cimento, vol 117B, No. 1, 15, (2002); J. Math. Phys., 42, 1682, (2001).

- Monin A.S. and Yaglon A.M. – “Statistical Fluid Mechanics” – Mit Press, Cambridge, vol. 2, 1971.
 - G. Rosen – Journ. Math. Phys., 12, 812, (1971).
 - Luiz C.L. Botelho, Mod. Phys. Lett 13B, 317, (1999).
 - A.A. Migdal, Int. Journal of Modern Physics A9, 1197, (1994).
 - V. Gurarie and A. Migdal, Phys. Rev. E54, 4908, (1996).
 - U. Frisch, “Turbulence” – Cambridge Press, Cambridge, (1996).
 - W. D. Mc-Comb – “The Physics of Fluid Turbulence, Oxford University, Oxford, (1990).
 - Luiz C.L. Botelho, Mod. Phys. Lett B13, 363, (1999).
 - Luiz C.L. Botelho, Nuov. Cimento 118B, 383, (2004)
 - Denisov SI, Horsthemke W., Phys. Lett A282 (6), 367, (2001).
 - Luiz C.L. Botelho, Int. J. Mod. Phys. B13, 1663, (1999).
 - Luiz C.L. Botelho, Mod. Phys. Lett B12, 301, (1998).
 - Cresson J.C.; Lyra M.L., J. Phys. Condens. Mat 8 (7), L83, (1996).
 - Luiz C.L. Botelho – Mod. Phys. Lett B12, 569, (1998).
 - Luiz C.L. Botelho – Mod. Phys. Lett B12, 1191, (1998).
 - Luiz C.L. Botelho – Mod. Phys. Lett B13, 203, (1999).
 - Luiz C.L. Botelho – Int. J. Mod. Phys. B12, 2857, (1998).
 - Luiz C.L. Botelho – Phys. Rev. 58E, 1141, (1998).
- [7] Michal Reed and Barry Simon, “Methods of Modern Mathematical Physics”, vol II, Academic Press, (1980), New York.